


Succinct Arguments


Lecture 06: Polynomial Commitments from Discrete Logarithms

Our recipe so far



Polynomial Commitments

Recall: Polynomial Commitments



- **Completeness:** Whenever $p(z) = v$, **R** accepts.
- **Extractability:** Whenever **R** accepts, **S**'s commitment **cm** “contains” a polynomial p of degree at most D .
- **Hiding:** **cm** and π reveal *no* information about p other than v

Cryptographic Groups

Group

A set $\mathbb{G}$ and an operation $*$

1. **Closure:** For all $a, b \in \mathbb{G}$, $a * b \in \mathbb{G}$
2. **Associativity:** For all $a, b, c \in \mathbb{G}$, $(a * b) * c = a * (b * c)$
3. **Identity:** There exists a unique element $e \in \mathbb{G}$ s.t. for every $a \in \mathbb{G}$,
 $e * a = a * e = a$.
4. **Inverse:** For each $a \in \mathbb{G}$, there exists $b \in \mathbb{G}$ s.t. $a * b = b * a = e$

E.g.: integers $\{ \dots, -2, -1, 0, 1, 2, \dots \}$ under $+$
positive integers mod prime $p : \{1, 2, \dots, p-1\}$ under $\times$
elliptic curves

Generator of a group

- An element g that generates all elements in the group by taking all powers of g

Examples: $\mathbb{F}_7^* := \{1,2,3,4,5,6\}$

$$\begin{array}{l} 3^1 = 3; \quad 3^2 = 2; \quad 3^3 = 6 \\ 3^4 = 4; \quad 3^5 = 5; \quad 3^6 = 1 \end{array} \pmod{7}$$

Discrete logarithm assumption

- A group $\mathbb{G}$ has an alternative representation as the powers of the generator $g: \{g, g^2, g^3, \dots, g^{p-1}\}$
- Discrete logarithm problem:
given $y \in \mathbb{G}$, find x s.t. $g^x = y$
- Example: Find x such that $3^x = 4 \bmod 7$
- Discrete-log assumption: discrete-log problem is computationally hard

Prime-order groups

- We will use only *prime-order groups*, i.e. groups where $|\mathbb{G}|$ is a large prime.
- Main examples of such groups are elliptic curve groups.
- We will call the field $\mathbb{F}_p$ the *scalar field* of the group.

Pedersen Commitment Scheme

Pedersen Commitments

$\text{Setup}(n \in \mathbb{N}) \rightarrow \text{ck}$

1. Sample random elements $g_1, \dots, g_n, h \leftarrow \mathbb{G}$

$\text{Commit}(\text{ck}, m \in \mathbb{F}_p^n; r \in \mathbb{F}_p) \rightarrow \text{cm}$

1. Output $\text{cm} := g_1^{m_1} g_2^{m_2} \dots g_n^{m_n} h^r$

Binding

Goal: For all efficient adv. $\mathcal{A}$,

$$\Pr \left[\text{Commit}(m; r) = \text{Commit}(m'; r') : \begin{array}{l} \text{ck} \leftarrow \text{Setup}(n) \\ (m, r, m', r') \leftarrow \mathcal{A}(\text{ck}) \end{array} \right] \approx 0$$

Proof: We will reduce to hardness of DL. Assume that $\mathcal{A}$ did indeed find breaking (m, r, m', r') . Let's construct $\mathcal{B}$ that breaks DL. Assume that $n = 1$.

Key idea: Let $h = g^x$. Then

$$g^m h^r = g^{m'} h^{r'} \implies g^{m+xr} = g^{m'+xr'}$$

$$\text{Can recover } x = \frac{m - m'}{r' - r}$$

- $\mathcal{B}(g, h)$

 1. $(m, r, m', r') \leftarrow \mathcal{A}(\text{ck} = (g, h))$
 2. Output $x = \frac{m - m'}{r' - r}$

Hiding

Goal: For all m, m' , and all adv. $\mathcal{A}$,
 $\mathcal{A}(\text{Commit}(m; r)) = \mathcal{A}(\text{Commit}(m'; r'))$

Proof idea: Basically one-time pad!

Let $\text{cm} := \text{Commit}(\text{ck}, m; r)$. Let $h = g^x$.

Then, for any m' , there exists r' such that $\text{cm} := \text{Commit}(\text{ck}, m'; r')$.

We could compute it, if we knew x : $r' = \frac{m - m'}{x} + r$

[Note: this doesn't break binding, because $\mathcal{A}$ doesn't know x

Additive Homomorphism

Let $\mathbf{cm}$ and $\mathbf{cm}'$ be commitments to m and m' wrt r and r' .
Then $\mathbf{cm} + \mathbf{cm}'$ is a commitment to $m + m'$ wrt $r + r'$

$$\begin{aligned}\mathbf{cm} &:= g_1^{m_1} \dots g_n^{m_n} h^r + \mathbf{cm}' := g_1^{m'_1} \dots g_n^{m'_n} h^{r'} \\ &= g_1^{m_1+m'_1} \dots g_n^{m_n+m'_n} h^{r+r'} \\ &= \text{Commit}(\text{ck}, m + m'; r + r')\end{aligned}$$

PC from DL-hard groups

PC scheme from Pedersen Comms

$\text{Setup}(d \in \mathbb{N}) \rightarrow (\text{ck}, \text{rk})$

1.

$\text{Commit}(\text{ck}, p \in \mathbb{F}_p^{d+1}; r \in \mathbb{F}_p) \rightarrow \text{cm}$

1.

$\text{Open}(\text{ck}, p, z \in \mathbb{F}_p; r) \rightarrow (\pi, v)$

1.

$\text{Check}(\text{rk}, \text{cm}, z, v, \pi) \rightarrow b \in \{0,1\}$

1.

PC scheme from Pedersen Comms

$\text{Setup}(d \in \mathbb{N}) \rightarrow (\text{ck}, \text{rk})$

1. $\text{ck} \leftarrow \text{Ped} . \text{Setup}(d + 1)$. Output $(\text{ck}, \text{rk}) = (\text{ck}, \text{ck})$.

$\text{Commit}(\text{ck}, p \in \mathbb{F}_p^{d+1}; r \in \mathbb{F}_p) \rightarrow \text{cm}$

- 1.

$\text{Open}(\text{ck}, p, z \in \mathbb{F}_p; r) \rightarrow (\pi, v)$

- 1.

$\text{Check}(\text{rk}, \text{cm}, z, v, \pi) \rightarrow b \in \{0, 1\}$

- 1.

PC scheme from Pedersen Comms

$\text{Setup}(d \in \mathbb{N}) \rightarrow (\text{ck}, \text{rk})$

1. $\text{ck} \leftarrow \text{Ped} . \text{Setup}(d + 1)$. Output $(\text{ck}, \text{rk}) = (\text{ck}, \text{ck})$.

$\text{Commit}(\text{ck}, p \in \mathbb{F}_p^{d+1}; r \in \mathbb{F}_p) \rightarrow \text{cm}$

1. Output $\text{cm} := \text{Ped} . \text{Commit}(\text{ck}, p; r)$

$\text{Open}(\text{ck}, p, z \in \mathbb{F}_p; r) \rightarrow (\pi, v)$

- 1.

$\text{Check}(\text{rk}, \text{cm}, z, v, \pi) \rightarrow b \in \{0, 1\}$

- 1.

PC scheme from Pedersen Comms

$\text{Setup}(d \in \mathbb{N}) \rightarrow (\text{ck}, \text{rk})$

1. $\text{ck} \leftarrow \text{Ped} . \text{Setup}(d + 1)$. Output $(\text{ck}, \text{rk}) = (\text{ck}, \text{ck})$.

$\text{Commit}(\text{ck}, p \in \mathbb{F}_p^{d+1}; r \in \mathbb{F}_p) \rightarrow \text{cm}$

1. Output $\text{cm} := \text{Ped} . \text{Commit}(\text{ck}, p; r)$

$\text{Open}(\text{ck}, p, z \in \mathbb{F}_p; r) \rightarrow (\pi, v)$

1. Output $(\pi := (p, r), v := p(z))$

$\text{Check}(\text{rk}, \text{cm}, z, v, \pi) \rightarrow b \in \{0, 1\}$

- 1.

PC scheme from Pedersen Comms

$\text{Setup}(d \in \mathbb{N}) \rightarrow (\text{ck}, \text{rk})$

1. $\text{ck} \leftarrow \text{Ped} . \text{Setup}(d + 1)$. Output $(\text{ck}, \text{rk}) = (\text{ck}, \text{ck})$.

$\text{Commit}(\text{ck}, p \in \mathbb{F}_p^{d+1}; r \in \mathbb{F}_p) \rightarrow \text{cm}$

1. Output $\text{cm} := \text{Ped} . \text{Commit}(\text{ck}, p; r)$

$\text{Open}(\text{ck}, p, z \in \mathbb{F}_p; r) \rightarrow (\pi, v)$

1. Output $(\pi := (p, r), v := p(z))$

$\text{Check}(\text{rk}, \text{cm}, z, v, \pi) \rightarrow b \in \{0, 1\}$

1. Check $\text{cm} = \text{Ped} . \text{Commit}(\text{ck}, p; r)$ and $p(z) = v$.

Completeness

Follows from correctness of Pedersen:
recomputing the commitment works.

Extractability

Follows from binding of Pedersen.

$\mathcal{E}(\mathbf{ck}, z)$

1. Invoke $\mathbf{cm} \leftarrow \mathcal{A}(\mathbf{ck})$
2. Get $(\pi = (p; r), v) \leftarrow \mathcal{A}(z)$.
3. Output p .

$\mathcal{E}$ outputs incorrect p if and only if $\mathcal{A}$ can provide a different opening for $\mathbf{cm}$

Hiding

Follows from hiding of Pedersen?

cm is perfectly hiding, but $\pi = (p, r)$ reveals polynomial!

Efficiency

$\mathbf{cm}$ is succinct (single $\mathbb{G}$ element), but $\pi = (p, r)$ is $O(d)$!

Better PC from DL

PC scheme from [BCGGP16]

Key idea: write polynomial as a $\sqrt{n} \times \sqrt{n}$ matrix, where n is num. coeffs

$$p = \begin{pmatrix} a_1 & \dots & a_m \\ a_{m+1} & \dots & a_{2m} \\ \vdots & & \\ a_{m(m-1)} & \dots & a_{m^2} \end{pmatrix}$$

Q: How to evaluate at z in matrix form?

$$p(z) = (1, z^m, \dots, z^{m(m-1)}) \begin{pmatrix} a_1 & \dots & a_m \\ a_{m+1} & \dots & a_{2m} \\ \vdots & & \\ a_{m(m-1)} & \dots & a_{m^2} \end{pmatrix} \begin{pmatrix} 1 \\ z \\ \vdots \\ z^{m-1} \end{pmatrix}$$

PC scheme from [BCGGP16]

$\text{Setup}(d \in \mathbb{N}) \rightarrow (\text{ck}, \text{rk})$

1. $\text{ck} \leftarrow \text{Ped}.\text{Setup}(\sqrt{d+1})$. Output $(\text{ck}, \text{rk}) = (\text{ck}, \text{ck})$.

$\text{Commit}(\text{ck}, p \in \mathbb{F}_p^{d+1}) \rightarrow \text{cm}$

- 1.

PC scheme from [BCGGP16]

$\text{Setup}(d \in \mathbb{N}) \rightarrow (\text{ck}, \text{rk})$

1. $\text{ck} \leftarrow \text{Ped} . \text{Setup}(d + 1)$. Output $(\text{ck}, \text{rk}) = (\text{ck}, \text{ck})$.

$\text{Commit}(\text{ck}, p \in \mathbb{F}_p^{d+1}) \rightarrow \text{cm}$

1. Write p as matrix $p = \begin{pmatrix} a_1 & \dots & a_m \\ a_{m+1} & \dots & a_{2m} \\ \vdots & & \\ a_{m(m-1)} & \dots & a_{m^2} \end{pmatrix}$
2. Use Pedersen to commit to rows, obtaining $\text{cm}_1, \dots, \text{cm}_m$
3. Output $\text{cm} := \begin{pmatrix} \text{cm}_1 \\ \vdots \\ \text{cm}_m \end{pmatrix}$

PC scheme from [BCGGP16]

$\text{Open}(\text{ck}, p, z \in \mathbb{F}_p; r) \rightarrow (\pi, v)$

1. Compute $\vec{z} := (1, z^m, \dots, z^{m(m-1)})$

2. Compute $\vec{a} = (1, z^m, \dots, z^{m(m-1)}) \begin{pmatrix} \uparrow & \dots & \uparrow \\ \vec{a}_1 & \dots & \vec{a}_m \\ \downarrow & \dots & \downarrow \end{pmatrix}$

3. Output $(\pi := \vec{a}, p(z))$

PC scheme from [BCGGP16]

$\text{Check}(\text{ck}, \text{cm}, z, v, \pi) \rightarrow b$

1. Parse $\text{cm} := \begin{pmatrix} \text{cm}_1 \\ \vdots \\ \text{cm}_m \end{pmatrix}$ and $\pi = (\vec{a})$
2. Compute $\vec{z} := (1, z^m, \dots, z^{m(m-1)})$
3. Compute $\text{pf} = (1, z^m, \dots, z^{m(m-1)}) \begin{pmatrix} \text{cm}_1 \\ \vdots \\ \text{cm}_m \end{pmatrix}$
4. Check $\text{pf} = \text{Ped} . \text{Commit}(\text{ck}, \vec{a})$
5. Check $v = \langle \vec{a}, (1, z, \dots, z^{m-1}) \rangle$

Completeness

Follows from homomorphism of Pedersen:

1. If $\mathbf{cm} := \begin{pmatrix} \text{cm}_1 = \text{Ped} . \text{Commit}(\text{ck}, \vec{a}_1) \\ \vdots \\ \text{cm}_m = \text{Ped} . \text{Commit}(\text{ck}, \vec{a}_m) \end{pmatrix}$
2. Then $\mathbf{pf} := (1, z^m, \dots, z^{m(m-1)}) \begin{pmatrix} \text{cm}_1 \\ \vdots \\ \text{cm}_m \end{pmatrix}$ commits to $\vec{a} = (1, z^m, \dots, z^{m(m-1)}) \begin{pmatrix} \vec{a}_1 \\ \vdots \\ \vec{a}_m \end{pmatrix}$
3. Additionally, by construction, $\vec{a}(z) = v$

Extractability

Follows from binding of Pedersen + rewinding

1. Extractor rewinds $\mathcal{A}$ n times, each time obtaining an evaluation at different points.
2. This gives us n linear equations in n unknowns, which we can solve.
3. Each iteration will be valid unless $\mathcal{A}$ breaks DL

Hiding

Follows from hiding of Pedersen?

cm is perfectly hiding, but $\pi = (\vec{a})$
reveals polynomial (but maybe less info?)

Efficiency

cm is $\sqrt{d}$ $\mathbb{G}$ elements, and π is $\sqrt{d}$ $\mathbb{F}_p$ elements.

Additionally, **Check** does only $O(\sqrt{d})$ work!